



# Developing a Theory-Based Intervention Manual to Enhance Self-Care of Patients with Heart Failure

Amanda Whittal, Vera Kalitzkus, Lou Atkins, Stefan Störk,  
Oliver Rudolf Herber (PI)

# Background

- International guidelines recommend self-care as integral part of routine heart failure (HF) management
- HF can be managed effectively with on-going self-care, yet patients are frequently unable to adhere
- Previous interventions that were not theory-based have shown limited success in improving adherence to self-care



# Aim of Study

- To develop an intervention manual containing theory-based BCIs that are well-defined using eight descriptors proposed to describe BCIs in a standardised way



# Research Question

- Can a detailed intervention manual for designing theory-based behaviour change interventions using the COM-B behaviour model improve self-care in HF patients?




# Study Design

- **Study design:** Use of COM-B model (Stage 1-3); Normalisation Process Theory (NPT) & Delphi technique (Stage 4) + Patient & Public Involvement
- **Duration:** 26 months (1FTE)
- **Value:** €220,114
- **Funding:** German Research Foundation (DFG)
- **Fund code:** DFG HE 7352/1-2
- **Ethical approval:** Ethics committee of HHU (Ref #: 2018-30)



# COM-B Model: Universal Behavioural Theory



# Stage 1: Extracting Behaviours

- Identification & extraction of *all* “target behaviours” associated with self-care (non-)adherence from two meta-studies (QUAN + QUAL)
- QUAL meta-summary (Herber *et al.* 2017) based on 31 reports
- QUAN meta-analysis (Kessing *et al.* 2016) based on 65 reports



# Stage 2: Mapping Behaviours onto COM-B

- Each of the factors identified in Stage 1 were mapped onto the COM-B model components (Capability, Opportunity, Motivation)
- If there were difficulties in classifying the factors onto the COM-B model, a second opinion was obtained
- The COM-B model assists in understanding of why patients with HF (non-)adhere to self-care





# Behaviour Change Wheel



# Stage 3: Identifying Behaviour Change Techniques

- Appropriate behaviour change techniques (BCTs) were identified for changing undesirable behaviours
- Use of Taxonomy (v1) that contains 93 BCTs



# Narrowing Determinant List: Less is More

- Merged target behaviours from QUAN + QUAL meta-studies
- Eliminated behaviours with effect sizes <25% (QUAL)
- Eliminated behaviours with unknown quality (QUAN)
- Combined overlapping determinants
- Focus on barriers only for larger intervention impact
- Enquired HF patients' preferences if several BCTs were available



# Role of Patient & Public Engagement

- 35 HF patients were asked to rate different BCTs in relation to its likelihood of use





# Next Steps

## Starting in November 2019



# Stage 4: Considering Contextual Factors

- Consultation of key stakeholders to identify wider factors needed for successful implementation of BCIs into routine work
- Qualitative semi-structured interviews with 15–17 key stakeholders (e.g. patients, nurses, doctors, researchers,...)
- Use of Normalisation Process Theory (NPT) provides guiding questions to overcome difficulties of implementing theoretically derived interventions into everyday practice



# Stage 4: Determination of Descriptors

- Interviews with key stakeholders will help determining the eight descriptors needed to describe BCIs in a standardised way





# Stage 4: Delphi Technique

- Use of Delphi technique (formal consensus method) involving *all* key stakeholders to elicit consensus on final interventions
- The Delphi questionnaire will deal specifically with any mixed responses (ambiguities) regarding the descriptors
- Threshold for consensus set at 75% of participating stakeholders; otherwise rank order will be used







# Contact Details

**Oliver Herber**

[Oliver.Herber@med.uni-duesseldorf.de](mailto:Oliver.Herber@med.uni-duesseldorf.de)

