

E
RESEARCH

The abstracts included in this brochure have been printed as submitted by the authors with some minor editing to maintain RCN house style. The RCN does not assume any responsibility for the accuracy or quality of the information provided in this book of abstracts. Authors are fully responsible for the content of their abstracts including accuracy of the facts, statements and references.

Publication code: 010 326

Contents

5

Monday 5 September
Theme 1.1 T eu Ct(C)10.8 (a)0.8 r( )]TJ0 Tc 0 Tw( )Tj78.837 0 Td( )Tj63.617 0 Td( )Tj63.617 0 Td( )T

Melbourne. She is Research and Education Lead for Nursing for the Victorian Comprehensive Cancer Centre (VCCC) Alliance, and co-lead of the Victorian Regional Research Teaching Hub. Mei is past President of the Clinical Oncology Society of Australia and the Cancer Nurses Society of Australia. Her research focuses on health equity and experiences of care for people with poor prognosis cancers, and the contribution of nurses to patient and system level outcomes.

Monday 5 September

2.15 –

azy 5 (e)1l32Tm( )T22-1..04 -0.7 (ac)9M4 -0.7 (M)Tj3(5)Tm( )TTc 0 Tw 2.06a

Dame Ruth May, Chief Nursing Officer, England

Biography

Professor Alex McMahon, Chief Nursing Officer for Scotland

Biography

Professor Alex McMahon is Chief Nursing Officer for Scotland and Director of the CNO Directorate in Scottish Government. He took up his post in January 2022
Previously he was Executive Director for Nursing, Midwifery and Allied Healthcare Professionals in NHS Lothian.

Professor McMahon has professional responsibility for two thirds of the NHS Scotland workforce: nursing, midwifery, AHPs and healthcare scientists. He leads on a number of policy areas such as infection prevention control.

Professor McMahon was the Executive Director of Nursing, Midwifery and AHPs at NHS Lothian and was Chair of the Scottish Executive Nurse Directors group for four years, working closely with the previous Chief Nursing Officer to shape policy and professional practice. He qualified as a registered mental health nurse in 1986, and as a registered general nurse in 1989, and has worked in the private sector, NHS and Scottish Government. He holds honorary professorships with the University of Stirling and Queen Margaret University.

Maria McIlgorm – Chief Nursing Officer, Department of Health, CNO Northern Ireland

Biography

Maria McIlgorm was appointed to the post of Chief Nursing Officer (CNO) for the Department of Health in March 2021. Prior to joining the Department, Maria was a professional advisor within the Scottish Government.

Maria joined the nursing register in 1988 before becoming a midwife in 1990. Since then,

Professor Hugh P. McKenna CBE, PhD, B.Sc.

and collaborated on studies examining the interconnections between mental health policy, the organisation and delivery of care, and the experiences of people using services.

individual bereavement follow up interventions or surveys the wider availability of services. This research evaluates the bereavement follow-up programme implemented in 2018 at the Queen Elizabeth University Hospital, Critical Care Unit.

Aims

- Evaluate the current intervention.
- Identify barriers and facilitators.
- Share the results within critical care.

Methods

This research is a mixed methods process evaluation following the UK MRC guidelines. (Moore, et al., 2014). Existing mortality and bereavement follow up data, and a relative's questionnaire examines the process variables through quantitative data. Qualitative interviews with key stakeholders and relatives engaging with the programme provide in-depth data about the experience of the intervention.

Results

94% of relatives want contact from critical care. Overall fidelity is affected by the reliance on a single contact and the collection of contact details. Engagement with the intervention is low, but the benefits for those who do return for a meeting go beyond answering clinical questions. Stakeholders want to support bereaved families but there are issues with organisational structure and resources. Influencing the results is the context of a large critical care department in the middle of a global pandemic.

Discussion and Conclusion

The results are mixed. Bereavement follow up is acceptable to relatives and stakeholders. Recognition of significant loss and the need for ongoing support are important mechanisms of change. However, there is a gap in relatives engaging with the programme due to a combination of implementation issues and a lack of understanding of the mechanisms of change. Changes to the programme are recommended.

Biography

Aileen Labram is currently a senior charge nurse in critical care at the Queen Elizabeth University Hospital in Glasgow. She is interested in bereavement support for families following the death of a loved one in critical care. Over the past two years she has carried out a research project to evaluate the bereavement follow-up programme introduced by herself and colleagues in critical care at the Queen Elizabeth University Hospital in 2018.

Research Topic: Acute an

Presenter: Louise Stayt, PhD, Oxford Brookes University, UK

Co-author(s): Clair Merriman, UK; Suzanne Bench, UK; Helen Walthall, UK; Sarah Vollum, UK; Jody Ede, UK; Ann Price, UK; Nicki Credland, UKKarin Gerber, UK; David Waters, UK

Abstract

Background

The emergence of COVID-19 created a surge in demand for critical care services leading to a large and sudden increase in critical care capacity through the reorganisation of beds and staff (Armocida et al., 2020; Wu & McGoogan, 2020). The specialist critical care nurse workforce was supplemented with staff re-deployed from other specialities which altered the way care was organised and delivered (Graselli et al. 2020). The impact of these changes on the quality and safety of patient care and the strategies employed by nurses

Biography

Louise is a senior lecturer in critical and specialist care at Oxford Brookes University. Louise's research interest include fatigue after critical illness, primary hypertension, patient safety, and fundamental nursing care. Louise is also a Professional Advisor for the British Association of Critical Care Nurses. Clair Merriman is a research active educationist and clinician. Clair currently holds the position as Principal Lecturer at Oxford Brookes

adults (over18s) undergoing cancer treatment. A narrative approach was used for analysis and integration of the findings. The analysis focused on behaviours conducted by carers to support patient's adherence across cancer sites and stages of treatment. The quality of studies was assessed in line with systematic review Advisory guidance. The systematic review protocol was registered in PROSPERO.

Results

Sixteen studies met our eligible criteria. Findings revealed three main themes, treatment responsibilities and decision-making (administering treatment, managing side-effects, and being involved in decisions); finding ways to meet treatment needs (seeking resources), and managing relationships and emotional reactions (communicating with patient/clinicians and providing emotional support).The changing role from being a spouse to acting like a .2 (m9.6 (s)-10.8 (u)-5M5)12 rieina (a).1 (g).0.7 (-13.2 2 T6 (vu.8 (i)12 (e3.2 f.6 82 f.6 912

research program in primary care nursing, with particular emphasis on nursing in general practice, chronic disease and nursing workforce issues.

Research Topic: Primary and Community Care

cases, the prevalence of missed care was more likely. Longer working hours in the community increased the risk of compromised care and sub-optimal patient care.

The aspects of the nursing process identified as ‘missed’ related to The World Health

Results

Participants were predominantly mothers (n=369/371, 99.4%), aged 25-34 (n=252, 67.8%), fit and healthy (n=314, 85%), white British (n=351, 94,5%) on maternity leave (n=252, 67.9%) and for half this was their first baby (186, 50.1%).

The lowest reported MCS was 29 (n=1), highest score 70 (n=4), mean score 59 (SD 6.5)d.e.7 (l)-10.4 (()-6

particularly during periods of hospitalisation which impacted their trust in healthcare professionals.

Conclusion

The findings illustrate the unique ways in which fathers made sense of their experiences of their child's care and how trust can be a fragile element of the parent / clinician relationship. The need for fathers to develop and sustain trust in healthcare professionals was a key finding of the study.

Biography

Peter is a Senior Lecturer in Children's Nursing at the School of Healthcare Sciences, Cardiff University. His clinical background is predominantly within paediatric critical care. His main areas of teaching include acute and critical care alongside clinical skills. Peter's PhD research utilised a narrative approach to explore fathers' experiences when their child is born with congenital heart disease. The research developed out of Peter's clinical experience, caring for children and young people with congenital heart disease. The research was funded by Cardiff University alongside a Florence Nightingale Foundation and Menat Trust research scholarship. Professor Daniel Kelly OBE was appointed RCN Chair of Nursing Research at Cardiff University in 2011. His main career focus has been in cancer care practice, management, research and education roles with an academic background in the social sciences at the University of Edinburgh and a PhD in Sociology at Goldsmiths, University of London. He is currently a Trustee at St Christopher's Hospice, and a Public Governor at Kings College Hospital NHS Foundation Trust. He is also Visiting Professor at UCL and The University of Edinburgh. His international work includes being Co-Chair of the HPV Action Network of the European Cancer Organisation which is Chair 12.2 (

Abstract

Background

Significant improvements in prognosis and treatment of childhood cancers (age 0-14) have increased the overall survival rates, with 85% now expected to survive more than five-years (National Health Service (NHS) 2022). These improvements are accompanied by prolonged and intensive treatments requiring extended time in health care settings and home-based treatments. It is inevitable that disruption created by cancer reaches beyond the diagnosed child, impacting the entire family. It is important to understand the experiences of siblings, to inform interventions to reduce the known associated distress

Research Topic: End of Life Care, Patient Experience, Inequalities in Health

Methodology: Other collection or analysis method

Research Approach: Other approaches

Presenter: Marie Clancy, RNC, Bachelor of Nursing, Masters in Public Health, PGCE in academic practice, Academy of Nursing, University of Exeter, UK

Abstract

Background

The challenge of researching sensitive and potentially vulnerable patient groups has been widely acknowledged in the literature. Whilst advisory group use is advocated by funding and ethical bodies, there is often little advice about tackling the challenges researchers may face when authentic engagement is desired. This presentation aims to highlight the importance of reflexivity to counter power imbalances and approach interpretations with cultural humilityffo i

Background

Research Topic: Older People

Methodology: Mixed

Research Approach:

Presenter: Helen Day, Professional Doctorate in Health and Social Care, South Tees Hospitals NHS Foundation Trust, UK

Abstract

Background

Concern around poor standards of nursing care for older people in hospital has been explored in relation to workload and operational pressures, but less around individuals' characteristics. Notions of personality traits and associated behaviours as possible influencers on nursing care delivery are variables that require consideration. The aim of this work is to critically explore behaviours of registered nursing staff working in older people's acute care settings from the perspectives of key stakeholders and identify whether there are any distinguishing personality traits that influence effective care delivery for older people.

Methodology

Only 34% and 32% reported enough knowledge to ask about possible UI and FI respectively. 40% did not where to refer people for continence symptoms within their organisation. Approximately half knew where to find in-house resources (52%) and online resources (47(i)1.1 31 (2.8t( )-11.7)-6.5 (p)1s p61-1.33781.774%ure5.5 (p)oited0.8 (p)7.7 9(u10.5 (c)10.4 (o)-

Co-author(s): Clare McKeaveney, UK; Claire Carswell, UK; Ann Bonner, Australia; Ilaria de Barbieri, Italy; William Johnston, UK

Biography

Joanne Reid is Professor of Cancer and Palliative Care and current chair of the Palliative Care Research Network, All Ireland Institute of Hospice and Palliative Care. She has conducted multi-disciplinary national and international funded studies and her publications reflect the international multi-disciplinary teams she leads (Joanne Reid – Research output — Queen's University Belfast (qub.ac.uk). Her dissemination has involved

Results

Nurses performed all 9 AMS behaviours. AMS behavioural performance was significantly higher ((238) = -4.14, = .000), by those who had training (= 53.15; SD = 7.40) compared to those who had not (= 48.30; = 10.75). All TDF determinants were also significantly higher in those with training. COVID-19 led to an increase in AMS behaviours, suggesting that nurses were extra vigilant.

Discussion

Standards of Proficienc

Abstract

Background
Missed nursing care is a phenomenon that compromises patient safety. Nurse managers may serve as facilitators for minimizing missed nursing care, by implementing certain strategies. However, there is lack of compilation of such strategies into formal recommendations, although a body of expertise exists among nurse managers. A special group of nurse managers is those working in nursing homes.

Aim
To identify strategies implemented by nurse managers working in nursing homes, on preventing and minimizing missed nursing care.

Methods
A qualitative study was conducted during 2019 and 2020, in four European countries (Italy, Germany, Switzerland, Cyprus), as part of the Nurse Managers for Safety Erasmus+ project. A purposive sample of nine nurse managers, working in nursing homes, participated in focus groups discussions and individual interviews. They described strategies and interventions on how they deal with the phenomenon of missed nursing care. Thematic analysis of the discussions and interviews and extraction of codes followed.

Results
Participants identified thirty-nine strategies grouped in seven themes, namely: i) empowerment and education, ii) improvement of care process iii) adequate staffing iv) communication and collaboration v) management of near missed incidents and adverse n ee (,)7.8 ( )0.8 (em)i.2 (ess.2 (ess.2(f)12.2 (n)i)111 (,)7.8 ( )-u)658tresin) v9.12 (r)1-1.8 ( )i.2(f)n-1.8 7 resif) em

video-conferencing. Interviews were fully transcribed, anonymised and analysed using Braun and Clarke's (2022) inductive, reflexive thematic approach to ensure key patterns and experiences were captured, relative to the study's research questions.

Results

Presenter: Owena Simpson, DAHP, LLM(LAMP), PGCEd, BSc(Hons), DipN, RGN., University of South Wales, UK

Abstract

There is currently a national nursing shortage which is driven by an ageing population and an increasing demand for health care. One initiative introduced to increase the number of registered nurses within the system has been to increase the number of students enrolled

2. Important content to be included within the onli

remove the ambiguity of clinical judgement when allocating an acuity level on an acute oncology assessment unit.

Aim

To determine accurate patient acuity on an acute oncology assessment unit, investigate trends and variations in acuity on admission, observe for correlation between acuity and admission source to inform safe staffing levels and future models of acute oncological care in the cancer centre.

Method

Retrospective analysis of acuity levels of recorded in real time on an acute oncology assessment unit from 31 January 2022 to 28 February 2022.

Results

There were 151 admissions to the acute oncology assessment unit between 8am and 8pm, the highest levels of acuity were seen between 5pm and 7pm.

Majority of the highest acuity patients presented unwell on site.

Research Approach: Evaluation (process, impact)

Presenter: Filip Haegdorens, RN, PhD, University of Antwerp, Belgium

Co-author(s): Carolien Wils, Belgium

Abstract

Background

Rapid response system activation is usually initiated by ward nurses with guidance of early warning systems such as the National Early Warning Score. However, when a patient is n 2 (n7 (v)1 (e)-1.)-0.9 2 (n04 70i6 (.)-1.W46.8 (-1.1 033 Td (e)-as (ar)n)-1. (at)-2.2) (al)10.p.2 271.1 -noh5as (ar)

Abstract

Background

Sexuality and intimacy are important to maintain quality of life, health and wellbeing among older adults (Jones, Moyle and Van Haitsma, 2021). These needs are often forgotten or ignored by health and social care professionals (HSCPs) in healthcare settings, particularly, for older people living with dementia and those identifying as lesbian, gay, bisexual, transgender, intersex or queer/questioning (LGBTIQ+) individuals (Horne et al.

sexuality via Delphi technique. She is interested in the issues related to sexuality, intimacy, ageing and dementia. She published a paper titled 'Exploratory study of Australian aged care staff knowledge and attitudes of later life sexuality' in 2017.

lacking, it negatively impacted upon Service Users' experiences of HSC. Supportive relationships within a community were instrumental in meeting the HSC needs of their individual members. Support that statutory services provided, did not always meet the needs of the people who were accessing or providing HSC, and they undervalued the support that communities provided.

Discussion

It is important to acknowledge the power of connection when designing HSC services that encourage supportive relationships between people who access and provide care, and their communities. Adopting Person-centred, Relationship based HSC could encourage meaningful connections and improve HSC experiences.

Conclusions

This study highlights indicators for improved sustainable HSC beyond COVID-19, advocating co-produced community-driven HSC services that meet the self-defined needs of those who access and provide them.

Biography

Louise is passionate about involving people and communities in co-produced Health and Social Care (HSC) and research. Her areas of interest include HSC education, community and voluntary sector wellbeing support, and project management. Louise completed her PhD study in December 2021, which explored the experiences and perceptions of people

Abstract

Background
The number of people living with dementia (PLWD) is predicted to rise from 55 million worldwide in 2017 to 139 million by 2050. PLWD in care homes are frequently denied their basic human rights and have reduced quality of life in comparison with those who live at home. There is wide disparity among care home services, both Nationally and Internationally. Staff working in care homes report high levels of stress and burnout. Access to dementia education varies.

Aims
This qualitative, ethically approved study aimed to understand and clarify the barriers and facilitators to upholding Human Rights for PLWD in care homes.

Method
In November 2019 seven focus groups were conducted. Each group consisted of 7-9 mixed participants, including staff from 20 nursing homes across Wales, PLWD, unpaid carers of PLWD and student nurses. Vignettes were used to base discussion around 'What are the of barriers and facilitators to upholding human rights for people living with dementia in care homes?'. Discussion was digitally recorded and transcribed verbatim. Thematic analysis was conducted using Braun and Clarke's (2006) method.

Results
Themes identified were barriers of **systems and processes**; including risk aversion, organisational culture, and financial constraints. **Personal barriers**: concerned values, beliefs, assumptions, knowledge, experience, and staff sense of worth. **Environmental barriers** included problems related to ‘false realities’, space, and privacy. **External pressures and expectations** from the public, family members and regulatory scrutiny were additional strains.

Discussion
Themes have informed the production of a bespoke e-learning resource for social care staff, and a further study to evaluate attitudes and understandings of care home staff in relation to dementia.

Conclusion
Safeguarding the human rights of PLWD in care homes is complex but must be prioritised. Overlapping barriers include organisational and personal factors, which need to be addressed on individual, cultural, societal and political levels.

Biography
Lesley is a Senior Lecturer in Nursing at Cardiff University. She has over 30 years’ experience in Nursing, most notably working in specialist roles focused on the care of older people in care homes, hospitals, and community settings. She has a Master’s degree in Psychoanalytic Psychotherapy. Lesley is passionate about applying and instilling person-centred care principles in caring for people living with Dementia. She teaches

extensively on this, across undergraduate and postgraduate programmes within Cardiff University, as well as delivering external presentations and webinars. She has conducted qualitative research on understanding the safeguarding of human rights for people living